

Biological nitrogen fixation by alternative nitrogenases in boreal cyanolichens: importance of molybdenum availability and implications for current biological nitrogen fixation estimates

Romain Darnajoux¹, Xinning Zhang², Darcy L. McRose², Jolanta Miadlikowska³, François Lutzoni³, Anne M. L. Kraepiel² and Jean-Philippe Bellenger¹

¹Centre Sève, Département de Chimie, Université de Sherbrooke, Sherbrooke, QC, Canada J1K 2R1; ²Department of Geosciences, Princeton University, Princeton, NJ 08544, USA;

³Department of Biology, Duke University, Durham, NC 27708, USA

Summary

Author for correspondence:

Jean-Philippe Bellenger

Tel: +1 819 821 7014

Email: jean-philippe.bellenger@usherbrooke.ca

Received: 16 May 2016

Accepted: 26 July 2016

New Phytologist (2017) **213**: 680–689

doi: 10.1111/nph.14166

Key words: alternative nitrogenases, biological nitrogen fixation, boreal forest, cyanolichens, molybdenum (Mo), *Peltigera aphthosa* s.l., vanadium (V).

• Cryptogamic species and their associated cyanobacteria have attracted the attention of biogeochemists because of their critical roles in the nitrogen cycle through symbiotic and asymbiotic biological fixation of nitrogen (BNF). BNF is mediated by the nitrogenase enzyme, which, in its most common form, requires molybdenum at its active site. Molybdenum has been reported as a limiting nutrient for BNF in many ecosystems, including tropical and temperate forests. Recent studies have suggested that alternative nitrogenases, which use vanadium or iron in place of molybdenum at their active site, might play a more prominent role in natural ecosystems than previously recognized.

• Here, we studied the occurrence of vanadium, the role of molybdenum availability on vanadium acquisition and the contribution of alternative nitrogenases to BNF in the ubiquitous cyanolichen *Peltigera aphthosa* s.l.

• We confirmed the use of the alternative vanadium-based nitrogenase in the *Nostoc* cyanobiont of these lichens and its substantial contribution to BNF in this organism. We also showed that the acquisition of vanadium is strongly regulated by the abundance of molybdenum.

• These findings show that alternative nitrogenase can no longer be neglected in natural ecosystems, particularly in molybdenum-limited habitats.

Introduction

The boreal biome covers 17% of the terrestrial landscape and represents one of the highest carbon storage pools in terrestrial ecosystems (Dixon *et al.*, 1994; IPCC, 2000; Kasischke, 2000; DeLuca & Boisvenue, 2012). Over the course of the next century, boreal forests will undergo changes at an unprecedented rate as a result of global climate change (IPCC, 2013, 2014). Nitrogen (N) is the most frequently reported limiting nutrient for primary production in boreal forests (LeBauer & Treseder, 2008; Wang *et al.*, 2010). The availability of N to plants is expected to play a critical role in the response of boreal forests to global climate change (Heimann & Reichstein, 2008; Sigurdsson *et al.*, 2013; Yuan & Chen, 2015). Biological nitrogen fixation (BNF) by cyanobacteria associated with bryophytes and lichen-forming fungi is a critical source of new N for unmanaged boreal ecosystems (DeLuca *et al.*, 2002).

Biological nitrogen fixation is catalyzed by the enzyme nitrogenase (Nase). Three iso-enzymes of Nase have been identified so far (Eady, 1996; Reed *et al.*, 2011): the canonical molybdenum (Mo)-Nase and two alternative Mo-independent isozymes, that

is, the vanadium (V)-dependent Nase (V-Nase) and the iron (Fe)-only Nase (Fe-Nase). Until very recently, the contribution of alternative Nases to N inputs in natural habitats was very difficult to measure, and BNF was thus usually assumed to be carried out exclusively by Mo-Nase. Yet, several studies carried out in tropical, temperate and cold temperate ecosystems have reported the limitation of BNF by Mo (Silvester, 1989; Barron *et al.*, 2008; Wurzbarger *et al.*, 2012; Jean *et al.*, 2013). V, which is *c.* 50–200 times more abundant than Mo in the Earth's crust (Wedepohl, 1995), might present a suitable alternative to Mo to sustain BNF in these environments. This is consistent with several studies reporting the presence of genes coding for alternative Nases in a wide range of ecosystems (Betancourt *et al.*, 2008) and in several cyanobacteria, including *Nostoc* spp. associated with five bi-membered (i.e. two symbionts) cyanolichens of the genus *Peltigera* (Hodkinson *et al.*, 2014). Another report, based on an exhaustive analysis of published acetylene reduction to ¹⁵N incorporation conversion ratios, concluded that both Mo-Nase and alternative Nases (V- and Fe-Nase) probably contribute to BNF in soils (Bellenger *et al.*, 2014). Finally, a recent study on metal homeostasis in the ubiquitous tri-membered (i.e. three

symbionts) cyanolichen *Peltigera aphthosa* s.l. also suggested that V probably plays a significant role in BNF by boreal cyanolichens (Darnajoux *et al.*, 2014).

Although these studies suggest that alternative Nases are widespread and active in a wide range of ecosystems, the definitive demonstration of their importance remains to be made. This goal is now within reach because new methods allow the assessment of the activity of alternative Nases in environmental samples. Recently, Zhang *et al.* (2014) have demonstrated that alternative N fixation produces organic matter with lower $\delta^{15}\text{N}$ (i.e. ^{15}N isotopic fractionation (ϵ) for Mo-Nase is *c.* +2‰, whereas, for alternative Nases, ϵ is *c.* +6 to +7‰). More recently, another isotope fractionation-based technique, isotopic acetylene reduction assay (ISARA), has been developed to efficiently and reliably assess alternative Nase activity in environmental samples (Zhang *et al.*, 2016).

Here, we make use of these new tools, together with more conventional approaches, to examine the contribution of alternative Nases to BNF in *P. aphthosa* s.l., and the environmental factors influencing this activity. *Peltigera aphthosa* s.l. is a symbiotic association between a fungus (*P. aphthosa*), a green alga (*Coccomyxa* sp.) and an N-fixing cyanobacterium (*Nostoc* sp.), confined to a specific structure called a cephalodium, which may contribute as much as 87% of potential BNF in boreal forest habitats in Sweden (Gavazov *et al.*, 2010; Rousk *et al.*, 2015). As only the gene for Mo-Nase and V-Nase has been found in *Nostoc* of *Peltigera* lichens to date (Hodkinson *et al.*, 2014) and, unlike V, Fe has a broad role in metabolism, we restricted our study to Mo- and V-Nase.

We hypothesized that the recently observed accumulation of V in *P. aphthosa* s.l. cephalodia (Darnajoux *et al.*, 2014) is widely distributed in the boreal biome. To test this hypothesis, we performed a discriminant analysis on the metal contents of *P. aphthosa* s.l. samples collected in several sites within the boreal biome. Considering the reported limitation of BNF by Mo and the known regulation of alternative Nases by Mo in many bacteria, we also hypothesized that Mo availability to cyanolichens is a critical environmental factor in the modulation of V acquisition and use for BNF in cyanolichens. This hypothesis was tested on a fire chronosequence in Sweden. As aerial depositions (e.g. particles, canopy flow-through) are the main sources of metals (e.g. V and Mo) to lichens (Darnajoux *et al.*, 2015; Marks *et al.*, 2015), we postulated that, along a vegetative succession, the amount of Mo and V, as well as the relative Mo : V ratio, might vary significantly as a result of different canopy openness and vegetation composition. A fire chronosequence might thus provide samples growing under similar climatic and geological conditions, but with contrasted Mo and V exposures. In addition, BNF has been shown to increase with stand age (Zackrisson *et al.*, 2004) on the selected fire chronosequence, offering a unique opportunity to determine whether or not Nase metal cofactor contents in cephalodia are influenced by stand age. Finally, we hypothesized that alternative Nases are present and actually contribute to BNF in *P. aphthosa* s.l. cyanolichens. This hypothesis was tested on *P. aphthosa* s.l. samples from the fire chronosequence using DNA, ISARA and $\delta^{15}\text{N}$ analyses.

Our results show that V is a metal of biological importance to *P. aphthosa* s.l. throughout the boreal biome, and provide direct

evidence confirming the hypothesis that V-Nase is present and contributes significantly to BNF by cyanolichens in boreal forest (Darnajoux *et al.*, 2014; Zhang *et al.*, 2016). We also demonstrate that Mo availability is a critical factor for the regulation of V inside the N-fixing symbiont of *P. aphthosa* s.l. Finally, we discuss to what extent the contribution of alternative Nases to BNF may affect the reliability of current BNF estimates reported in the literature.

Materials and Methods

Sampling

Specimens of *Peltigera aphthosa* (L.) Willd. s.l. were collected in Alberta (Wood Buffalo National Park, August 2013, $n=6$) and from a 369-yr-old fire chronosequence in Sweden (September 2013, $n=16$, details on the fire chronosequence can be found in Zackrisson *et al.*, 2004). Data from samples in northern Québec, Alaska (Fairbanks), Russia (Stolby Reserve, Krasnoyarsk) and Fjord du Saguenay (FdS, South Québec) ($n=27$, 5, 6 and 9, respectively; Darnajoux *et al.*, 2014) were also used in the statistical analyses. Additional samples (S1–S6, individual thalli) were collected for the ISARA experiment in August 2014 in four sites along the Swedish fire chronosequence (S1, 115 yr; S2, 282 yr; S3, 304 yr; S4–S6, 369 yr). All thalli were manually cleaned of foreign materials (e.g. bryophytes and leaves) with forceps and then kept dry in opaque paper bags at room temperature until further processing.

Thallus preparation and elemental analysis

Lichens were washed to remove metals present at the surface of the thallus (nonspecific absorption sites and particles). Algae and cephalodia were separated from the thallus of *P. aphthosa* s.l. and elemental contents of unwashed thallus, washed thallus, algae and cephalodia were determined by inductively coupled plasma-mass spectrometry (ICP-MS), as described in Darnajoux *et al.* (2014). Metal contents are reported as either mol^{-1} phosphorus ($\text{mol}_{\text{metal}} \text{mol}_{\text{P}}^{-1}$) or g^{-1} of thallus biomass ($\mu\text{g}_{\text{metal}} \text{g}_{\text{thallus}}^{-1}$).

$\delta^{15}\text{N}$ experiments

MilliQ-washed ground lichen thalli were analyzed for $\delta^{15}\text{N}$ at the 'Laboratoire d'analyses des isotopes stables et légers' at GEOTOP-UQAM. Data are expressed as ‰ vs air ($\pm 0.2\text{‰}$ for 1σ).

Isotopic acetylene reduction assay

Acetylene was produced by reacting CaC_2 (Acros Organic, Springfield, NJ, USA) with water and collecting the resulting gas in a Tedlar bag. Samples from Sweden (S1–S6), collected in 2014, 2 months before analysis, were sprayed with deionized water and then left to acclimate at 10°C in an incubator (Infors HT multitron pro, Anjou, QC, Canada) with a 12 h : 12 h, light : dark cycle for at least 5 d, with regular spraying (every 2 d)

to maintain optimal humidity. Samples were then incubated in 250-ml Mason jars with 10% headspace removed and replaced with acetylene. Lichens were left for 1–6 d under a 12 h : 12 h, light : dark cycle at 10°C, with regular sampling of the headspace. Ethylene production and background concentration of ethylene in each acetylene batch were determined by gas chromatography as described in Jean *et al.* (2013). Linearity was confirmed up to 90 h in all samples. All gas samples were then sent for ISARA analyses at Princeton University (for more information on the method, see Zhang *et al.* (2016)).

Results are presented as the ^{13}C fractionation of the reduction of acetylene to ethylene ($^{13}\epsilon_{\text{AR}} (\text{‰}) = \delta^{13}\text{C}_{\text{acetylene}} - \delta^{13}\text{C}_{\text{ethylene}}$, at low ethylene yield). The fraction of alternative fixation in acetylene reduction assay (ARA) was determined assuming a linear model from $^{13}\epsilon_{\text{AR},\text{Mo}}$ and $^{13}\epsilon_{\text{AR},\text{V}}$, and the percentage of alternative BNF activity was calculated as the fraction of alternative fixation in ARA corrected with $\text{C}_2\text{H}_4 : \text{N}_2$ R values ($R_{\text{Mo}} = 3$, $R_{\text{V}} = 1$). Calculation details can be found in Supporting Information Methods S1 and in Zhang *et al.* (2016).

DNA extractions and PCR amplifications

DNA was extracted from dissected cephalodia of samples S1–S6 using the QuickExtract Plant DNA Extraction Solution (Epicentre, Madison, WI, USA), according to the manufacturer's instructions (65°C for 6 min, 98°C for 2 min, chill at 4°C). DNA extracts were diluted 1 : 25 in nuclease-free water in order to avoid PCR inhibition. Degenerate *vnfD* PCR primers were designed to correspond roughly to the region amplified by Bellenger *et al.* (2014) with bases adapted to *Nostoc vnfD* sequences from Hodkinson *et al.* (2014). Primer sequences were as follows: F (5'-TYAACATGG GGTGGATGA-3'); R (5'-CCATTGTGGTAGCCATG-3'). PCRs of 25 µl were conducted using 2U Expand High Fidelity DNA Polymerase (Roche Applied Science, Indianapolis, IN, USA) per reaction, 200 µM deoxynucleoside triphosphates (dNTPs), 500 nM forward and reverse primers and 2 µl of DNA template per reaction. PCR conditions for all reactions were as follows: 94°C for 2 min, followed by 38 cycles of 94°C for 15 s, 50°C for 30 s and 72°C for 2 min, with a final elongation at 72°C for 7 min. PCR products were gel purified using the QIAquick gel extraction kit (Qiagen, Valencia, CA, USA), ligated into the TOPO 4.0 vector (Invitrogen, Life Technologies, Grand Island, NY, USA) and cloned into One-Shot TOP10 (Invitrogen, Life Technologies) chemically competent *Escherichia coli*. Plasmid DNA was purified with the QIAprep Miniprep kit (Qiagen) and unidirectional Sanger DNA sequencing was conducted at Macrogen (Macrogen USA, New York, NY, USA) using the M13F(-47) primer (5'-CGCCAGG GTTTTCCAGTCACGAC-3'). Eight and five *vnfD* sequences were analyzed from specimens collected at S1 (GenBank accession number: KX502814) and S4 (GenBank accession number: KX502813), respectively.

Statistical analyses

Data for regression and multivariate discriminate analyses were log-transformed in order to avoid spurious correlation or to

achieve normality. Canonical discriminant analysis (CDA) was performed using R software v.3.1.2 (RSTUDIO 0.98.501, package FACTOMINEr 1.24, CANDISC 0.6-5 and NORTEST 1.0-2; R Development Core Team, 2009). All regressions were performed with R using a linear model (LM and GLM, STAT 3.1.2), with selection of variables based on the significance of regression ANOVA (function ANOVA, CAR 2.0-22) and on the minimization of the Akaike information criterion (AIC). All residuals were tested for normality. Figures were drawn using either R or SIGMAPLOT 11.2.0.5 (Systat Software, San Jose, CA, USA), and were edited using INKSCAPE v.0.48.4 (Freeware, <http://inkscape.org>).

Results

Discriminant analysis on the metal contents in the symbiotic partners of *P. aphthosa* s.l. across all sampling sites

To evaluate the biological relevance of V for boreal cyanolichens, that is *P. aphthosa* s.l. (Darnajoux *et al.*, 2014), at a broad scale, we performed a discriminant analysis on metal contents of 63 specimens of *P. aphthosa* s.l. collected in boreal forests in pristine areas (northern Alberta and Québec), a low contamination area (Sweden) and higher contamination areas (Alaska and Russia) (Fig. S1). Metal contents were analyzed in four different ways; unwashed thallus, washed thallus, algal layer and cephalodium. The discriminant analysis was able to correctly separate all compartments (error rate, 0.031; Table S1). The first canonical axis mostly separates unwashed thalli from cephalodia, whereas the second canonical axis separates the algal layer from everything else (Fig. 1). The washed thallus group has a central position in the canonical score plot. Arrows indicate the loading of variables on the canonical axes; the longer an arrow towards a group, the higher its participation to discriminate this group from others. The analysis identified manganese (Mn), zinc (Zn) and lead (Pb) as discriminants for unwashed thallus; magnesium (Mg) and titanium (Ti) as discriminant for algae; and Fe, Mo, V, chromium (Cr), copper (Cu) and cobalt (Co) as discriminant for cephalodia (Fig. 1). Similar results were obtained for samples from contaminated and uncontaminated sites (compare Fig. 1a and b).

Relationship between atmospheric deposition and Mo and V contents in lichen cephalodia across pristine and low metal contamination sites

In order to evaluate the impact of atmospheric deposition on metal contents in the cephalodia, we analyzed Mo and V contents in unwashed thalli (in $\mu\text{g}_{\text{metal}} \text{g}_{\text{thallus}}^{-1}$) and cephalodia (in $\text{mol}_{\text{metal}} \text{mol}_{\text{P}}^{-1}$) in samples collected in pristine and low metal contamination sites ($n = 58$). Metal contents in unwashed thalli are assumed to be closely correlated with atmospheric deposition fluxes (Darnajoux *et al.*, 2015). Samples from Alaska and Russia were not considered as they are characterized by higher metal deposition that can interfere with natural metal management (homeostasis) (Fig. S1). As shown in Fig. 2(a), Mo cellular quotas in cephalodia are positively correlated with Mo exposures (i.e.




Fig. 1 Metals in tri-membered lichen *Peltigera aphthosa* s.l. Canonical (Can) discriminant analysis of all thalli of *P. aphthosa* s.l. ($n = 63$) with regard to compartments in (a) all samples and (b) nonheavily contaminated boreal sites (Alberta, North Québec and Sweden) (see Supporting Information Fig. S1a). UWT, unwashed thalli (black triangles); WT, oxalate–EDTA-washed thalli (brown circles); ALG, algae (green crosses); CEP, cephalodia (blue plus symbols). All data in $\text{mol}_{\text{metal}} \text{mol}_{\text{P}}^{-1}$ were log-transformed before the analysis. Data from Fjord du Saguenay (South Québec) were not included as algal data were not available.

Mo content in unwashed thalli, $P < 0.001$; details on regression analyses can be found in Table S2). In contrast with Mo, V cellular quotas in cephalodia are relatively constant in all samples and are not correlated with V exposure (Fig. 2b), a clear indication that V content in cephalodia, the site of N fixation, is tightly regulated. Further analysis of the data revealed that Mo and V cellular quotas are correlated in cephalodia (Fig. 2c; quadratic correlation, $P = 0.003$; Table S1), with Mo and V being positively correlated until Mo concentrations reach $(1-2) \times 10^{-4} \text{ mol}_{\text{Mo}} \text{mol}_{\text{P}}^{-1}$, and negatively correlated afterwards.

Mo and V contents in *P. aphthosa* s.l. samples collected along a fire chronosequence in Sweden

We analyzed Mo and V contents in unwashed thallus, washed thallus, algal layer and cephalodium of *P. aphthosa* s.l. samples collected at five sites along a fire chronosequence (Fig. 3). In unwashed and washed thalli, Mo is present at lower concentrations than V. Mo and V covary across sites (Fig. 3a,b), with higher concentrations at Reivo, Tjadnes and Vaksliden (50, 282 and 304 yr) than at Guorbåive and Ruttjeheden (185 and 369 yr). In the algal layer, Mo and V have similar concentrations and do not vary significantly across sites (Fig. 3c). In the cephalodia, V and Mo concentrations are of the same order, but are clearly anticorrelated (Fig. 3d). Further analysis of the data revealed that the V concentration is positively correlated with the Mo concentration in unwashed thallus, but is negatively correlated with the Mo concentration in cephalodia (Fig. 3e; Table S2).

ISARA measurement on *P. aphthosa* s.l. samples from Sweden

We used ISARA to determine the contribution of V-Nase to BNF on freshly collected *P. aphthosa* s.l. samples from northern Sweden (S1–S6), characterized by low to moderate metal depositions (Fig. S1a). The results show that four of the six *Peltigera* samples tested have a lower $^{13}\epsilon_{\text{AR}}$ value than that expected for Mo-only Nase activity (Fig. 4a). The estimated percentages of alternative BNF (Methods S1) show that alternative Nases contribute to BNF (mean \pm SD = $35 \pm 12\%$; range, 20–60%) in all samples (Fig. 4b; see later Table 2). The presence of *vnfD* genes was confirmed in samples S1 (GenBank accession number

KX502814) and S4 (GenBank accession number KX502813). BLASTN searches of GenBank (April 2015) showed highest identity to sequences from lichens deposited by (Hodkinson *et al.*, 2014; accessions: KF662359, KF662360, KF662361, KF662362).

Characterization of $\delta^{15}\text{N}$ in *P. aphthosa* s.l. samples from Sweden

We determined the $\delta^{15}\text{N}$ values for 14 of the 16 thalli of *P. aphthosa* s.l. from Fig. 3 (Fig. 5). The $\delta^{15}\text{N}$ averaged value ranged from 0.8 to 1.7‰ (individual values ranged from -1.4 to 3.2‰), with the highest averaged value measured in Tjadnes and the lowest averaged value measured in Guorbåive (individual extremes are found in Ruttjeheden). Reivo, Vaksliden and Ruttjeheden achieved intermediate values (Fig. 5a). The comparison of the $\delta^{15}\text{N}$ and V concentration in the cephalodia showed a negative correlation between these two datasets (Fig. 5b), with lower $\delta^{15}\text{N}$ corresponding to higher V : P ratios in the cephalodia.

Discussion

Occurrence of V in *P. aphthosa* s.l. cephalodia

Discriminant analysis performed on *P. aphthosa* s.l., collected along a natural gradient of metal deposition (Alberta, Quebec, Sweden, Alaska and Russia, Fig. S1a), identified Fe, Mo and V as discriminators for cephalodia (Fig. 1). This is consistent with Nases being important Fe and Mo reservoirs in N_2 fixers (Kustka *et al.*, 2003; Bellenger *et al.*, 2011). This also suggests that the presence of V in cyanolichen cephalodia is widely distributed in the boreal biome, from pristine to contaminated areas.

Mo and V accumulation in *P. aphthosa* s.l. cephalodia as a function of deposition

Considering that atmospheric deposition is an important source of metal to lichens, we evaluated whether or not the presence of Mo and V in cephalodia reflects atmospheric deposition. The strong correlation between Mo concentrations in cephalodia and atmospheric Mo deposition (Fig. 2a), in pristine and low metal contamination sites, suggests that Mo deposition primarily




Fig. 2 Linear regression of molybdenum (Mo) and vanadium (V) in cephalodia (CEP) of *Peltigera aphthosa* s.l. thalli from nonheavily contaminated boreal sites (northern Alberta, Québec and Sweden). All data were log-transformed (axes are in \log_{10} scale) to achieve normality of residues and to avoid spurious correlation. (a) Mo : phosphorus (P) ($\text{mol}_{\text{Mo}} \text{mol}_{\text{P}}^{-1}$) in cephalodia as a function of Mo exposure ($\mu\text{g}_{\text{Mo}} \text{g}_{\text{thallus}}^{-1}$). (b) V : P ($\text{mol}_{\text{V}} \text{mol}_{\text{P}}^{-1}$) in cephalodia as a function of V exposure ($\mu\text{g}_{\text{V}} \text{g}_{\text{thallus}}^{-1}$). (c) V : P ($\text{mol}_{\text{V}} \text{mol}_{\text{P}}^{-1}$) in cephalodia as a function of Mo : P ($\text{mol}_{\text{Mo}} \text{mol}_{\text{P}}^{-1}$) in cephalodia. $n = 58$. Short dashed lines show 95% prediction intervals and long dashed lines show 95% confidence intervals.

controls Mo availability to cephalodia. In contrast with Mo, V cellular quotas in cephalodia are not correlated with V deposition (Fig. 2b; Table S2). Instead, V cellular quotas in cephalodia in all samples are kept at a fairly constant level (2.0 ± 0.3) $\times 10^{-4} \text{ mol}_{\text{V}} \text{mol}_{\text{P}}^{-1}$, which is close to the optimum V cellular quotas (V : P) required to sustain diazotrophic growth through V-Nase in cyanobacteria ($(2-5) \times 10^{-4} \text{ mol}_{\text{V}} \text{mol}_{\text{P}}^{-1}$; Darnajoux *et al.*, 2014). This suggests a tight regulation of V content in

cephalodia independent of V exposure, possibly related to BNF. The correlation between V and Mo cellular quotas in cephalodia (quadratic correlation, $P = 0.003$; Table S2) suggests that Mo availability might control V accumulation in cephalodia. For Mo cellular quotas ($\text{Mo} : \text{P}) < (1-2) \times 10^{-4} \text{ mol}_{\text{Mo}} \text{mol}_{\text{P}}^{-1}$, Mo and V are positively correlated in the cephalodia, reflecting their correlation in metal exposure (Fig. S1b). This trend is observed primarily among the samples collected in the pristine areas of Québec and Alberta. We hypothesize that metal deposition in these sites is so low that cyanolichens use both Mo and V to sustain BNF. For Mo cellular quotas in cephalodia of $> (1-2) \times 10^{-4} \text{ mol}_{\text{Mo}} \text{mol}_{\text{P}}^{-1}$, Mo and V are negatively correlated, suggesting that Mo availability regulates, or at least impacts, V accumulation in cephalodia.

Role of Mo on V accumulation in *P. aphthosa* s.l. cephalodia

In order to further investigate the role of Mo availability on V accumulation in cephalodia, we focused on *P. aphthosa* s.l. samples collected along a fire chronosequence in Sweden. Samples collected along this fire chronosequence show a range of Mo and V deposition, as illustrated by the unwashed and washed thalli data (Fig. 3a,b). Contrary to our initial hypothesis, Mo and V concentrations in the thalli do not show systematic variations with stand age. Variation in Mo and V exposure appears to be site specific. The covariation of Mo and V concentration in unwashed thalli suggests common sources for both Mo and V all along the chronosequence. The higher abundance of V compared with Mo in unwashed thalli reflects the relative abundance of Mo and V in the environment. The Mo content in cephalodia followed the same trend as metal deposition, confirming that Mo deposition strongly influences Mo availability to cephalodia (compare Fig. 3a and d). However, V cellular quotas in cephalodia do not reflect V deposition. Instead, V cellular quotas in cephalodia follow an opposite trend when compared with Mo, and are low when Mo deposition and Mo in the cephalodia are high (Fig. 3d). Data from the fire chronosequence (local metal gradient) are consistent with data from metal surveys across pristine and contaminated sites (large-scale metal gradient), and further support our hypothesis that V accumulation in cephalodia is primarily controlled by Mo abundance in cephalodia (Fig. 3e; Table S2), rather than V deposition. This regulation of V by Mo is not observed in the algal symbiont (Fig. 3c), suggesting that it is related to BNF in the cephalodia.

Our data suggest that the presence of V in the cephalodia of *P. aphthosa* s.l. is ubiquitous in the boreal biome and that Mo availability is an important factor controlling V accumulation. This is consistent with the hypothesis that alternative Nases can support BNF in Mo-limited ecosystems. The presence of V in the cephalodia (minimum of $c. 4 \times 10^{-5} \text{ mol}_{\text{V}} \text{mol}_{\text{P}}^{-1}$), even when Mo is abundant, suggests that V can play a role in BNF, even when Mo is not limiting. The effect of temperature comes to mind, as experiments on purified enzymes have demonstrated that V-Nase is less affected than Mo-Nase by low temperature




Fig. 3 Nitrogenase (Nase) metal cofactors in cyanolichens collected along a 369-yr fire chronosequence. Molybdenum (Mo) and vanadium (V) contents in (a) unwashed thalli (UWT), (b) washed thalli (WT), (c) green algae (ALG) and (d) cephalodia (CEP). (e) Linear regression of Mo : phosphorus (P) ratio vs V : P ratio in cephalodia (CEP) and unwashed thalli (UWT). Data were log-transformed to avoid spurious correlation. Error bars are \pm SE; $n = 3, 2, 3, 3, 5$ for the 50-yr-old (Reivo burn, Rvo), 185-yr-old (Guorbåive, Gbv), 282-yr-old (Tjadnes, Tjd), 304-yr-old (Vaksliden, Vkd) and 369-yr-old (Ruttjeheden, Rjd) sites, respectively.




Fig. 4 Nitrogenase (Nase) utilization in *Peltigera aphthosa* s.l. collected in Sweden based on isotopic acetylene reduction assay (ISARA). (a) ¹³ε_{AR} data from pure culture of various diazotrophs for molybdenum (Mo)-Nase only (closed circles), vanadium (V)-Nase only (white circles) and from *P. aphthosa* s.l. (gray circles, samples S1–S6). Gray box represents the range of values obtained for Mo-Nase (13.1–14.7‰). Error bars represent estimated \pm SE for a single ethylene measurement. (b) Estimated proportions of biological fixation of nitrogen (BNF) by alternative Nases for *P. aphthosa* s.l. (more details on the calculations can be found in Zhang *et al.*, 2016, and in Methods S1). Samples are individual thalli collected along a 369-yr fire chronosequence: S1 (115 yr), S2 (282 yr), S3 (304 yr), S4–S6 (369 yr). Error bars were calculated using error propagation.

(Miller & Eady, 1988). In addition, at low temperature (<14°C), the expression of V-Nase genes is no longer repressed by Mo (Walmsley & Kennedy, 1991). Thus, the low temperature in

boreal ecosystems is likely to favor alternative Nases and might explain, in part, the occurrence of V in cephalodia, even at high Mo content.




Fig. 5 Nitrogen isotopic content in *Peltigera aphthosa* s.l. collected along a 369-yr fire chronosequence. (a) $\delta^{15}\text{N}$ of washed thallus (WT); (b) regression (solid line) of $\delta^{15}\text{N}$ in washed thallus vs vanadium (V) content in cephalodia (CEP, $\text{mol}_{\text{metal}} \text{mol}_{\text{P}}^{-1}$). Data are averaged for each fire chronosequence site. Error bars are \pm SE; $n = 3, 2, 3, 3, 5$ for the 50-yr-old (Reivo burn, Rvo), 185-yr-old (Guorbåive, Gbv), 282-yr-old (Tjadnes, Tjd), 304-yr-old (Vaksliiden, Vdk) and 369-yr-old (Ruttjeheden, Rjd) sites, respectively. Short dashed lines show 95% prediction intervals and long dashed lines show 95% confidence intervals.

Contribution of alternative Nases to BNF by *P. aphthosa* s.l.

The presence of V in cephalodia and its strong correlation with Mo availability suggest an active contribution of this alternative Nase to BNF by *P. aphthosa* s.l. As Mo availability significantly impacts V acquisition, we further evaluated whether or not Mo:V abundances in cephalodia (Fig. 3d) reflect the contribution of alternative Nases to BNF in the field using $\delta^{15}\text{N}$ data, which provide an integrated view of the various N sources over the lifetime of the lichen. As reported by Zhang *et al.* (2014), N fractionation can be used to evaluate the contribution of alternative Nases ($\delta^{15}\text{N}$ c. -6 to 8%) vs Mo-Nase ($\delta^{15}\text{N}$ c. -2%) in organic matter.

Variations in $\delta^{15}\text{N}$ in *P. aphthosa* s.l. from the fire chronosequence sites in Sweden are consistent with the metal content measured in cephalodia (Fig. 5a). $\delta^{15}\text{N}$ decreases significantly when V concentration in the cephalodia increases (Fig. 5b; $P = 0.023$; Table S2), which is in accordance with a higher contribution of alternative Nases to BNF. Although these data must be interpreted with caution, as other factors, such as N deposition, could also contribute to variations in the $\delta^{15}\text{N}$ data, they suggest that alternative Nases might contribute to a greater extent to BNF in Mo-limited sites.

The presence of *vnfD* genes in samples from the Swedish chronosequence was established in two of the six samples tested (S1 and S4), corroborating a previous publication on the presence of alternative Nases in cyanolichens from the genus *Peltigera* (Hodkinson *et al.*, 2014). In addition, ISARA data collected on a subset of lichen samples from the Swedish chronosequence showed that alternative Nases contribute significantly to BNF (mean \pm SD = $35 \pm 12\%$; range, 20–60%) in laboratory incubations at low temperature (10°C) (Fig. 4).

ISARA results and $\delta^{15}\text{N}$ data are consistent and suggest that alternative Nases are not only active in short-term laboratory experiments (ISARA data), but are also likely to contribute to BNF by these cyanolichens in the field ($\delta^{15}\text{N}$ data), especially in sites characterized by low Mo availability and low temperatures. Considering that most cyanolichens are physiologically active

during cold periods of the year (MacFarlane & Kershaw, 1977; Muir *et al.*, 1997), further research is required to fully characterize the effect of temperature on the relative contribution of alternative Nases to BNF by boreal cyanolichens in Mo-rich and Mo-poor ecosystems.

Re-evaluation of BNF estimates

Our results, together with others (Zhang *et al.*, 2016), imply that alternative Nases contribute to BNF in several boreal forest ecosystems, which could significantly change our understanding of the metal requirements of, and enzymes involved in, BNF. These findings also raise questions regarding the reliability of current BNF estimates. The canonical method used to assess BNF in field samples, that is, ARA (Hardy *et al.*, 1968), requires the use of a conversion ratio (ARA:N₂) in order to convert ARA data into BNF (N inputs). This ratio can be affected by many factors, such as growth conditions, but primarily by the type of Nase in use; ARA:N₂ = 3:1, 1–2:1 and <1:1 for the Mo-, V- and Fe-only Nases, respectively (Bellenger *et al.*, 2014). However, because the calibration of ARA data by ^{15}N incorporation experiments is problematic for many environmental samples, the

Table 1 Summary of literature survey (Supporting Information Table S3) on biological nitrogen fixation by cryptogamic cover (N₂-fixing bacteria associated with bryophytes and lichens)

	No. of studies	Acetylene reduction assay-based studies	^{15}N calibration made	Percentage of ^{15}N calibration
Arctic	2	1	0	0
Boreal	11	10	6	55
Continental	6	5	1	17
Temperate	12	12	6	50
Tropical	7	6	3	43
Dry	7	7	1	14
Total	45	41	17	38

Table 2 Estimation of nitrogen fixation ($\mu\text{g N g}_{\text{lichen}}^{-1} \text{d}^{-1}$) capacity of *Peltigera aphthosa* s.l. when considering molybdenum (Mo)-only contribution ($\text{C}_2\text{H}_4 : ^{15}\text{N}_{\text{Mo}} = 3 : 1$) or estimation of activity of alternative Nases with isotopic acetylene reduction assay ($\text{C}_2\text{H}_4 : ^{15}\text{N}_{\text{alt}} = 1 : 1$)

Sample no.	C_2H_4 reduction ($\mu\text{mol C}_2\text{H}_4 \text{ g}_{\text{lichen}}^{-1} \text{d}^{-1}$)	N fixation Mo-only ($\mu\text{g N g}_{\text{lichen}}^{-1} \text{d}^{-1}$)	N fixation ISARA ($\mu\text{g N g}_{\text{lichen}}^{-1} \text{d}^{-1}$)	Percentage error/ ISARA estimation
1	1.6	15	22	46
2	3.5	32	40	23
3	3.4	31	41	30
4	1.7	16	18	16
5	2.2	21	26	26
6	2.1	20	27	34

Percentage error represents the difference between estimates based on $\text{C}_2\text{H}_4 : ^{15}\text{N}_{\text{Mo}} = 3 : 1$ and $\text{C}_2\text{H}_4 : ^{15}\text{N}_{\text{alt}} = 1 : 1$.

theoretical Mo-Nase conversion ratio of $\text{ARA} : \text{N}_2 = 3$ is often assumed. A survey of > 40 reports on BNF by cryptogamic cover, which could possibly account for nearly one-half of BNF on unmanaged land (Elbert *et al.*, 2012), highlights that $\text{ARA} : ^{15}\text{N}$ conversion ratios were effectively measured in less than one-half of the studies (Tables 1, S3). Another potential bias comes from the experimental determination of $\text{ARA} : \text{N}_2$ ratios. As a result of low BNF activity, ^{15}N incorporation data are often acquired under laboratory conditions that are poorly representative of the field (i.e. increased temperature, CO_2 and humidity, Table S3). Recent reports on the presence of easily absorbed forms of ^{15}N in commercial $^{15}\text{N}_2$ gas stocks (Dabundo *et al.*, 2014) further emphasize the importance of interpreting past reports on BNF with great caution.

Interestingly, studies that have performed both ARA and ^{15}N incubations in the field on the same samples have revealed that soil, lichen and moss samples often yield intermediate ratios, indicating simultaneous N fixation by Mo-Nase and alternative Nases (Menge & Hedin, 2009; Bellenger *et al.*, 2014; Vile *et al.*, 2014). Thus, current BNF estimates probably include large errors, and the magnitude of these errors remains to be fully determined. However, the assumption of the predominance of Mo-Nase, without appropriate validations, could result in a significant underestimation of N inputs (16–46%, Table 2). Considering that alternative Nases have been found virtually everywhere in terrestrial ecosystems, with the exception of symbiotic associations with higher plants, it is an urgent matter to better characterize the enzymatic diversity of BNF in various compartments of the boreal forest and other ecosystems in order to re-evaluate BNF estimates. One crucial step is to deploy more rigorous protocols, allowing scientists to properly convert field measurements of Nase activity into N input estimates by implementing, for instance, new methods and tools (such as ISARA and $\delta^{15}\text{N}$), leading to better evaluations of the contribution of alternative Nases to BNF.

Conclusions

Our data confirm that alternative Nases are not only present in boreal cyanolichens, but are also active and probably contribute to BNF by cyanolichens throughout the boreal biome. Although Mo availability and temperature seem to be important factors controlling Nase activity, significant work remains to be performed to fully understand the conditions under which

alternative Nases come into play. It is also crucial to develop and implement methods, such as a combination of ARA, ISARA and $\delta^{15}\text{N}$, to measure BNF and the relative contributions of Mo and alternative Nases in field samples. A better characterization of Nase activity, including alternative Nases, in boreal forests and other ecosystems, will significantly improve our understanding of the role of micronutrients on ecosystem function, improve our estimates of BNF and may help to clarify N budgets.

Acknowledgements

We wish to thank Mikhail Zhurbenko, as well as Anastasya Knorre and Altya Dutbaeva, both at 'Stolby' National Wildlife Nature Reserve (Krasnoyarsk, Siberia, Russia), for the field assistance in Russia; Air Saguenay pilot Jacques Bérubé and the team at Lac Margane for the sampling in northern Québec, Canada, as well as the Société des établissements de plein air du Québec (SEPAQ) Fjord du Saguenay; and members of the Lutzoni laboratory (Bernie Ball, Emilie Lefevre) and the Arnold laboratory (University of Arizona) (A. Elizabeth Arnold and Kayla Arendt) for the field work in Québec. We also thank Marie-Charlotte Nilsson-Hegethorn and members of her research team for their support in sample collection in Sweden. The sampling of lichen specimens for this project was, in part, performed with the financial support of the National Science Foundation through a Dimensions of Biodiversity award DEB-1046065 to F.L. and a REVSYS award on the genus *Peltigera* DEB-1025930 to J.M. and F.L. This work was supported by the Canadian Research Chair in terrestrial biogeochemistry (CRC-950-219383), the Natural Sciences and Engineering Research Council of Canada through a Discovery grant (RGPIN-386963-2011) and the Fond Québécois de Recherche, Nature et Technologie through a Nouveau chercheur grant (FRQNT-2014NC-172330).

Author contributions

J-P.B. and R.D. conceived the study. A.M.L.K. and X.Z. designed the ISARA experiments. F.L., J.M., J-P.B. and R.D. collected lichen specimens. F.L. and J.M. identified the lichens. R.D., X.Z. and D.L.M. performed the experiments and analyzed the data. R.D. wrote the first draft of the manuscript. All authors contributed to the final version of the manuscript.

References

- Barron AR, Wurzburger N, Bellenger J-P, Wright SJ, Kraepiel AML, Hedin LO. 2008. Molybdenum limitation of asymbiotic nitrogen fixation in tropical forest soils. *Nature Geoscience* 2: 42–45.
- Bellenger J-P, Wichard T, Xu Y, Kraepiel AML. 2011. Essential metals for nitrogen fixation in a free-living N₂-fixing bacterium: chelation, homeostasis and high use efficiency. *Environmental Microbiology* 13: 1395–1411.
- Bellenger J-P, Xu Y, Zhang X, Morel FMM, Kraepiel AML. 2014. Possible contribution of alternative nitrogenases to nitrogen fixation by asymbiotic N₂-fixing bacteria in soils. *Soil Biology and Biochemistry* 69: 413–420.
- Betancourt DA, Loveless TM, Brown JW, Bishop PE. 2008. Characterization of diazotrophs containing Mo-independent nitrogenases, isolated from diverse natural environments. *Applied and Environmental Microbiology* 74: 3471–3480.
- Dabundo R, Lehmann MF, Treibergs L, Tobias CR, Altabet MA, Moisaner PH, Granger J. 2014. The contamination of commercial ¹⁵N₂ gas stocks with ¹⁵N-labeled nitrate and ammonium and consequences for nitrogen fixation measurements. *PLoS ONE* 9: e110335.
- Darnajoux R, Constantin J, Miadlikowska J, Lutzoni F, Bellenger J-P. 2014. Is vanadium a biometal for boreal cyanolichens? *New Phytologist* 202: 765–771.
- Darnajoux R, Lutzoni F, Miadlikowska J, Bellenger J-P. 2015. Determination of elemental baseline using peligeralean lichens from Northeastern Canada (Québec): initial data collection for long term monitoring of the impact of global climate change on boreal and subarctic areas in Canada. *Science of the Total Environment* 533: 1–7.
- DeLuca TH, Boisvenue C. 2012. Boreal forest soil carbon: distribution, function and modelling. *Forestry* 85: 161–184.
- DeLuca TH, Zackrisson O, Nilsson M-C, Sellstedt A. 2002. Quantifying nitrogen-fixation in feather moss carpets of boreal forests. *Nature* 419: 917–920.
- Dixon RK, Solomon AM, Brown S, Houghton RA, Trexler MC, Wisniewski J. 1994. Carbon pools and flux of global forest ecosystems. *Science* 263: 185–190.
- Eady RR. 1996. Structure–function relationships of alternative nitrogenases. *Chemical Reviews* 96: 3013–3030.
- Elbert W, Weber B, Burrows S, Steinkamp J, Büdel B, Andreae MO, Pöschl U. 2012. Contribution of cryptogamic covers to the global cycles of carbon and nitrogen. *Nature Geoscience* 5: 459–462.
- Gavazov KS, Soudzilovskaia NA, van Logtestijn RSP, Braster M, Cornelissen JHC. 2010. Isotopic analysis of cyanobacterial nitrogen fixation associated with subarctic lichen and bryophyte species. *Plant and Soil* 333: 507–517.
- Hardy RWF, Holsten RD, Jackson EK, Burns RC. 1968. The acetylene–ethylene assay for N₂ fixation: laboratory and field evaluation. *Plant Physiology* 43: 1185–1207.
- Heimann M, Reichstein M. 2008. Terrestrial ecosystem carbon dynamics and climate feedbacks. *Nature* 451: 289–292.
- Hodkinson BP, Allen JL, Forrest LL, Goffinet B, Sérusiaux E, Andrésón ÓS, Miao V, Bellenger J-P, Lutzoni F. 2014. Lichen-symbiotic cyanobacteria associated with *Peltigera* have an alternative vanadium-dependent nitrogen fixation system. *European Journal of Phycology* 49: 11–19.
- IPCC. 2000. Watson RT, Noble IR, Bolin B, Ravindranath NH, Verardo DJ, Dokken DJ, eds. *IPCC special report: land use, land-use change, and forestry. Summary for policymakers*. Geneva, Switzerland: Cambridge University Press.
- IPCC. 2013. Stocker TF, Qin D, Plattner G-K, Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V, Midgley PM, eds. *Climate change 2013: the physical science basis. Contribution of Working Group I to the fifth assessment report of the Intergovernmental Panel on Climate Change*. Cambridge, UK & New York, NY, USA: Cambridge University Press.
- IPCC. 2014. Core Writing Team, Pachauri RK, Meyer LA, eds. *Climate change 2014: synthesis report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Geneva, Switzerland: IPCC.
- Jean ME, Phalyvong K, Forest-Drolet J, Bellenger J-P. 2013. Molybdenum and phosphorus limitation of asymbiotic nitrogen fixation in forests of eastern Canada: influence of vegetative cover and seasonal variability. *Soil Biology and Biochemistry* 67: 140–146.
- Kasischke ES. 2000. Boreal ecosystems in the global carbon cycle. In: Kasischke ES, Stocks BJ, eds. *Fire, climate change, and carbon cycling in the boreal forest*. New York, NY, USA: Springer-Verlag, 19–30.
- Kustka A, Carpenter EJ, Capone DG, Raven JA. 2003. A revised estimate of the iron use efficiency of nitrogen fixation, with special reference to the marine cyanobacterium *Trichodesmium* spp. (Cyanophyta). *Journal of Phycology* 39: 12–25.
- LeBauer D, Treseder K. 2008. Nitrogen limitation of net primary productivity. *Ecology* 89: 371–379.
- MacFarlane JD, Kershaw KA. 1977. Physiological–environmental interaction in lichen. IV. Seasonal changes in the nitrogenase activity of *Peltigera canina* (L.) Willd var. *Praetextata* (Floerke in Somm.) Hue, and *P. canina* (L.) Willd var. *Rufescens* (Weiss) Mudd. *New Phytologist* 79: 403–408.
- Marks JA, Pett-Ridge JC, Perakis SS, Allen JL, McCune B. 2015. Response of the nitrogen-fixing lichen *Lobaria pulmonaria* to phosphorus, molybdenum, and vanadium. *Ecosphere* 6: 1–17.
- Menge DNL, Hedin LO. 2009. Nitrogen fixation in different biogeochemical niches along a 120 000-year chronosequence in New Zealand. *Ecology* 90: 2190–2201.
- Miller RW, Eady RR. 1988. Molybdenum and vanadium nitrogenases of *Azotobacter chroococcum*. Low temperature favors N₂ reduction by vanadium nitrogenase. *Biochemical Journal* 256: 429–432.
- Muir PS, Shirazi AM, Patrie J. 1997. Seasonal growth dynamics in the lichen *Lobaria pulmonaria*. *The Bryologist* 100: 458–464.
- R Development Core Team. 2009. *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Reed SC, Cleveland CC, Townsend AR. 2011. Functional ecology of free-living nitrogen fixation: a contemporary perspective. *Annual Review of Ecology, Evolution, and Systematics* 42: 489–512.
- Rousk K, Sorensen PL, Lett S, Michelsen A. 2015. Across-habitat comparison of diazotroph activity in the subarctic. *Microbial Ecology* 69: 778–787.
- Sigurdsson BD, Medhurst JL, Wallin G, Eggertsson O, Linder S. 2013. Growth of mature boreal Norway spruce was not affected by elevated [CO₂] and/or air temperature unless nutrient availability was improved. *Tree Physiology* 33: 1192–1205.
- Silvester W. 1989. Molybdenum limitation of asymbiotic nitrogen fixation in forests of Pacific Northwest America. *Soil Biology and Biochemistry* 21: 283–289.
- Vile MA, Kelman Wieder R, Živković T, Scott KD, Vitt DH, Hartsock JA, Iosue CL, Quinn JC, Petix M, Fillingim HM et al. 2014. N₂-fixation by methanotrophs sustains carbon and nitrogen accumulation in pristine peatlands. *Biogeochemistry* 121: 317–328.
- Walmsley J, Kennedy C. 1991. Temperature-dependent regulation by molybdenum and vanadium of expression of the structural genes encoding three nitrogenases in *Azotobacter vinelandii*. *Applied and Environmental Microbiology* 57: 622–624.
- Wang YP, Law RM, Pak B. 2010. A global model of carbon, nitrogen and phosphorus cycles for the terrestrial biosphere. *Biogeosciences* 7: 2261–2282.
- Wedepohl HK. 1995. The composition of the continental crust. *Geochimica et Cosmochimica Acta* 59: 1217–1232.
- Wurzbürger N, Bellenger JP, Kraepiel AML, Hedin LO. 2012. Molybdenum and phosphorus interact to constrain asymbiotic nitrogen fixation in tropical forests. *PLoS One* 7: 2–8.
- Yuan ZY, Chen HYH. 2015. Decoupling of nitrogen and phosphorus in terrestrial plants associated with global changes. *Nature Climate Change* 5: 1–5.
- Zackrisson O, DeLuca TH, Nilsson M-C, Sellstedt A, Berglund LM. 2004. Nitrogen fixation increases with successional age in boreal forests. *Ecology* 85: 3327–3334.
- Zhang X, McRose DL, Darnajoux R, Bellenger J-P, Morel FMM, Kraepiel AML. 2016. Alternative nitrogenase activity in the environment and nitrogen cycle implications. *Biogeochemistry* 127: 189–198.
- Zhang X, Sigman DM, Morel FMM, Kraepiel AML. 2014. Nitrogen isotope fractionation by alternative nitrogenases and past ocean anoxia. *Proceedings of the National Academy of Sciences, USA* 111: 4782–4787.

Supporting Information

Additional Supporting Information may be found online in the Supporting Information tab for this article:

Fig. S1 Unwashed thallus and cephalodia contents of vanadium (V) and molybdenum (Mo).

Table S1 Discrimination of lichen compartment with respect to metals

Table S2 Regression parameters for linear regression in Figs 2, 3 (e) and 5(b)

Table S3 Literature survey of biological nitrogen fixation (BNF) estimation and measurement conditions for di-nitrogen-fixing species in various ecosystems around the world

Methods S1 Contribution of alternative nitrogenase to acetylene reduction and N₂ fixation.

Please note: Wiley Blackwell are not responsible for the content or functionality of any Supporting Information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.



About *New Phytologist*

- *New Phytologist* is an electronic (online-only) journal owned by the New Phytologist Trust, a **not-for-profit organization** dedicated to the promotion of plant science, facilitating projects from symposia to free access for our Tansley reviews.
- Regular papers, Letters, Research reviews, Rapid reports and both Modelling/Theory and Methods papers are encouraged. We are committed to rapid processing, from online submission through to publication 'as ready' via *Early View* – our average time to decision is <28 days. There are **no page or colour charges** and a PDF version will be provided for each article.
- The journal is available online at Wiley Online Library. Visit **www.newphytologist.com** to search the articles and register for table of contents email alerts.
- If you have any questions, do get in touch with Central Office (np-centraloffice@lancaster.ac.uk) or, if it is more convenient, our USA Office (np-usaoffice@lancaster.ac.uk)
- For submission instructions, subscription and all the latest information visit **www.newphytologist.com**